

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 639 574 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: 94112652.6

(51) Int. Cl.<sup>6</sup>: **C07D 417/12, C07D 277/56,  
A01N 43/78**

(22) Date of filing: 12.08.94

(30) Priority: 16.08.93 KR 9315846

(43) Date of publication of application:  
22.02.95 Bulletin 95/08(84) Designated Contracting States:  
**BE CH DE DK FR GB IT LI NL**(71) Applicant: **LUCKY LTD.**  
20, Yoido-dong,  
Yongdungpo-ku  
Seoul 150-721 (KR)

(72) Inventor: **Rew, Yo Sup**  
Lucky Dormitory 206,  
386-1, Doryong-dong  
Youseong-ku,  
Daejeon (KR)  
Inventor: **Cho, Jin Ho**  
Lucky Hana Apt. 101-502,  
Shinseong-dong  
Youseong-ku,  
Daejeon (KR)  
Inventor: **Ra, Choon Sup**  
Woobang Apt. 101-303,  
138-6, Jungpyong-dong  
Kyongsan-shi,  
Kyongbuk (KR)  
Inventor: **Ahn, Sei Chang**  
Lucky Apt. 6-405,  
381-42, Doryong-dong  
Youseong-ku,  
Daejeon (KR)  
Inventor: **Kim, Sung Kee**  
Lucky Apt. 9-303,  
381-42, Doryong-dong  
Youseong-ku,


**Daejeon (KR)**  
Inventor: **Lee, Yong Hwan**  
Lucky Apt. 6-206,  
381-42, Doryong-dong  
Youseong-ku,  
**Daejeon (KR)**  
Inventor: **Jung, Bon Young**  
Chukong Apt. 106-502,  
Beop-dong  
Deaduk-ku,  
**Daejeon (KR)**  
Inventor: **Choi, Woo Bong**  
Lucky Hana Apt. 109-1201,  
Shinseong-dong  
Youseong-ku,  
**Daejeon (KR)**  
Inventor: **Rhee, Young Ho**  
Chukong Apt. 210-1008,  
Beop-Dong  
Daeduk-ku,  
**Daejeon (KR)**  
Inventor: **Yoon, Man Young**  
Lucky Dormitory 409,  
386-1, Doryong-dong  
Youseong-ku,  
**Daejeon (KR)**  
Inventor: **Chun, Sung Wook**  
Lucky Dormitory 311,  
386-1, Doryong-dong  
Youseong-ku,  
**Daejeon (KR)**

(74) Representative: **Fürniss, Peter, Dr.**  
**Kuhnen, Wacker & Partner**  
Alois-Steinecker-Strasse 22  
D-85354 Freising (DE)

(54) 2-aminothiazolecarboxamide derivatives, processes for their preparation and their use for controlling phytopathogenic organisms.


(57) The present invention relates to a novel 2-aminothiazolecarboxamide derivative represented by the following general formula (I):

**EP 0 639 574 A1**



in which

R<sup>1</sup> and R<sup>2</sup> independently of one another represent hydrogen, (C<sub>1</sub>-C<sub>5</sub>)alkyl, (C<sub>1</sub>-C<sub>5</sub>)haloalkyl, (C<sub>3</sub>-C<sub>6</sub>)-alkenyl, (C<sub>3</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,



R<sup>3</sup> or phenyl or benzyl, each of which can be substituted with halogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl or nitro;

represents (C<sub>1</sub>-C<sub>3</sub>)alkyl or (C<sub>1</sub>-C<sub>3</sub>)haloalkyl;

R<sup>4</sup> represents 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, phenyl, or phenyl or benzyl, each of which can be substituted with halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or nitro;

R<sup>5</sup> represents cyano or thiocarbamoyl; and

R<sup>6</sup> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)alkenyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, or phenyl or benzyl, each of which can be substituted with halogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl or nitro,


which is a fungicidal agent useful for controlling phytopathogenic organisms.

In addition, the present invention also relates to a process for preparing the novel 2-aminothiazolecarboxamide derivative of formula (I) and use of the compound of formula (I) as an agent for controlling phytopathogenic organisms.


## BACKGROUND OF THE INVENTION

## 1. Field of the invention

5 The present invention relates to a novel thiazolecarboxamide derivative. More particularly, the present invention relates to a novel 2-aminothiazolecarboxamide derivative represented by the following general formula (I):



15 in which  
 R<sup>1</sup> and R<sup>2</sup> independently of one another represent hydrogen, (C<sub>1</sub>-C<sub>5</sub>)alkyl, (C<sub>1</sub>-C<sub>5</sub>)haloalkyl, (C<sub>3</sub>-C<sub>6</sub>)alkenyl, (C<sub>3</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,




25 or phenyl or benzyl, each of which can be substituted with halogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl or nitro;  
 R<sup>3</sup> represents (C<sub>1</sub>-C<sub>3</sub>)alkyl or (C<sub>1</sub>-C<sub>3</sub>)haloalkyl;  
 R<sup>4</sup> represents 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, phenyl, or phenyl or benzyl, each of which can be substituted with halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or nitro;  
 30 R<sup>5</sup> represents cyano or thiocarbamoyl; and  
 R<sup>6</sup> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)alkenyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, or phenyl or benzyl, each of which can be substituted with halogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl or nitro.

The present invention also relates to a process for preparing the compound of formula (I), as defined above, and use of the compound of formula (I) for controlling phytopathogenic organisms.

## 2. Background Art


35 Fungicidal agents against Oomycetes occupy approximately 25% of the total fungicide market and has built up a unique market because Oomycetes is characterized by mycelium having no ergosterol biosynthesis mechanism. At the present time, conventional non-systemic fungicidal agents, for example, captan,  
 40 captafol, dithiocarbamate, chlorothalonil, etc., have been widely used for controlling Oomycetes. However, it has been disclosed that such non-systemic fungicidal agents have only a preventive activity against Oomycetes but has a serious toxic effect. In the 1970's acylalanine-based fungicidal agents have been developed as a systemic fungicide. However, it has been also reported that the resistance of Oomycetes to  
 45 such systemic fungicides becomes gradually aggravated. Further, recently the field of fungicidal agents against Oomycetes does not achieve any remarkable development. Accordingly, the development of a novel fungicide against Oomycetes which has a high fungicidal activity and a low toxicity has been strongly desired.

To satisfy such requirement, in the latter half of the 1980's dimethomorph was developed and reported  
 50 as an agent for controlling causative organisms of late blight and downy mildew in fruit trees and potatoes. In addition, for disinfecting cereal seeds and controlling causative organisms of rust and smut metsulfosax represented by the following formula (II) was developed (see S. African Patent No. 67 06,681) :



Thereafter, the following thiazolecarboxamide-based compounds have also been proposed as an effective substance having fungicidal activity against Oomycetes :

10 1) Heterocyclic carboxamide compounds represented by the following general formula (see European Patent No. 268,892) :




20 in which

A denotes pyrazolyl, imidazolyl, thiazolyl, isothiazolyl, oxazolyl or isoxazolyl, each of which has substituent(s),

B denotes alkoxy, alkylthio, imidazolyl, pyrazolyl, furyl or thienyl, and


D denotes cyano, thiocarbamoyl or acylthiocarbamoyl.

25 Among the above-identified heterocyclic carboxamide compounds, the compound having the following formula (III) has been proposed as a typical effective substance :



and has been reported as having a significant preventive and curative effects on *Pseudoperonospora cubensis*, which is the causative organism of downey mildew in cucumber, at the concentration of 100ppm.

40 2) Thiazolecarboxamide compounds represented by the following general formula (see European Patent No. 292,937 and Japanese Laid-open Patent Publication No. (Hei) 4-154704 (1992)) :




in which

R<sup>1</sup> and R<sup>2</sup> represent hydrogen or (C<sub>1</sub>-C<sub>3</sub>)alkyl, and


R<sup>3</sup> represents 2-furyl, 3-furyl, 2-thienyl or 3-thienyl.

55 Among the above-identified thiazolecarboxamide compounds, the compound having the following formula (IV) has been proposed as a typical effective substance :



10 and it has been reported that the compounds (IV-1) and (IV-2) prevent tomato late blight by 70% and 80%, respectively, at the concentration of 25ppm.

3) Thiazolecarboxamide compounds represented by the following general formula (see European Patent No. 313,091) :



20 in which


one of X and Y represents S and the other represents C,

Z represents cyano or thiocarbamoyl,

25 R<sup>1</sup> and R<sup>2</sup> independently of one another represent hydrogen, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halomethyl or phenyl, and

R<sup>3</sup> represents (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)haloalkenyl, furyl, thienyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)-alkylthio, (C<sub>3</sub>-C<sub>5</sub>)alkynyloxy, (C<sub>3</sub>-C<sub>5</sub>)alkynylthio, pyrazolyl, or phenyl which can be substituted with halogen.

4) Thiazolecarboxamide compounds represented by the following general formula (see European Patent No. 434,620) :



40 in which


R<sup>1</sup> and R<sup>2</sup> independently of one another represent hydrogen, alkyl, cycloalkyl (which can be substituted with methyl or methylthio) or -CH<sub>2</sub>XR<sup>4</sup>,

R<sup>3</sup> represents furyl or thienyl,

45 R<sup>4</sup> represents alkyl, alkenyl or alkynyl (each of which can be substituted with halogen or alkoxy), or phenyl or benzyl (each of which can be substituted with halogen, alkyl, alkoxy, trifluoromethyl or nitro), and

X represents oxygen or sulfur atom.

50 As the typical effective substance of the above-identified compounds, the compound having the following formula (V) :



10 was proposed and it has been reported that the compound of formula (V) controls tomato late blight by 90% or more at the concentration of 60ppm.

The above exemplified thiazolecarboxamide effective compounds have a certain degree of fungicidal activity and also a low toxicity. However, they have a disadvantage in that their fungicidal activities are lower than that of dimethomorph which was developed and commercialized in the latter half of the 1980's.

15 Thus, the present inventors have extensively studied to develop a new compound having a high fungicidal activity and a low toxicity. As a result, we have identified that a certain thiazolecarboxamide derivative having an amino group attached to 2-position of thiazole ring, i.e. a novel 2-aminothiazolecarboxamide derivative having the general formula (I), as defined above, shows a potent fungicidal activity which is higher than those of the presently known thiazolecarboxamide compounds and dimethomorph, and also has a systemic activity and a curative activity as well as a preventive activity, and then completed the present invention.

Therefore, it is an object of the present invention to provide a novel 2-aminothiazolecarboxamide derivative having the general formula (I), as defined above.


25 It is a further object of the present invention to provide a process for preparing the 2-aminothiazolecarboxamide derivative of formula (I).

It is another object of the present invention to provide use of the 2-aminothiazolecarboxamide derivative of formula (I) as an agent for controlling plant diseases caused by typical phytopathogenic organisms of Oomycetes.

30 The more pertinent and important features of the present invention have been outlined above in order that the detailed description of the invention which follows will be better understood and that the present contribution to the art can be fully appreciated. Those skilled in the art can appreciate that the conception and the specific embodiment disclosed herein may be readily utilized as a basis for modifying or designing other structures for carrying out the same purpose of the present invention. Further, those skilled in the art can realize that such equivalent constructions do not depart from the spirit and scope of the invention as set forth in the claims.


## DISCLOSURE OF INVENTION

40 In one aspect, the present invention relates to a novel 2-aminothiazolecarboxamide derivative represented by the following general formula (I) :



in which

50 R<sup>1</sup> and R<sup>2</sup> independently of one another represent hydrogen, (C<sub>1</sub>-C<sub>5</sub>)alkyl, (C<sub>1</sub>-C<sub>5</sub>)haloalkyl, (C<sub>3</sub>-C<sub>6</sub>)alkenyl, (C<sub>3</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,



or phenyl or benzyl, each of which can be substituted with halogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl or nitro;

R<sup>3</sup> represents (C<sub>1</sub>-C<sub>3</sub>)alkyl or (C<sub>1</sub>-C<sub>3</sub>)haloalkyl;

R<sup>4</sup> represents 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, phenyl, or phenyl or benzyl, each of which can be substituted with halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or nitro;

R<sup>5</sup> represents cyano or thiocarbamoyl; and

5 R<sup>6</sup> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)alkenyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, or phenyl or benzyl, each of which can be substituted with halogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl or nitro.

The compound of formula (I) of the present invention has a potential fungicidal activity against typical phytopathogenic organisms of Oomycetes including Pythiaceae, Peronosporaceae, etc., for example, Plasmopara viticola in grapes, Phytophthora infestans in potato and tomato, Phytophthora capsici in red  
10 pepper, and the like, and therefore, can be used as an agent for controlling plant diseases caused by such phytopathogenic organisms.

Among the compound of for formula (I), the preferred one includes those wherein R<sup>1</sup> and R<sup>2</sup> independently of one another represent hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)alkenyl, (C<sub>3</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl or R<sup>3</sup> represents methyl, ethyl or trifluoromethyl, R<sup>4</sup> represents 2-thienyl or 3-thienyl, R<sup>5</sup>  
15 represents cyano or thiocarbamoyl and R<sup>6</sup> represents (C<sub>1</sub>-C<sub>6</sub>)alkyl.

The compound of formula (I) according to the present invention can also exist in the form of an optically active isomer. Therefore, it should be understood that such optical isomer of the compound of formula (I) is included within the scope of the present invention.

In addition, the present invention also relates to a process for preparing the novel 2-aminothiazolecarboxamide derivative having the general formula (I) above. The process of the present invention can be  
20 represented by the following synthetic methods (I) and (II).

25

30

35



40

45


50

55

Synthetic Method (I)




(I)  $\text{R}^5$ =cyano group





(I)

$\text{R}^5$ =thiocarbamoyl group

Synthetic Method (II)



X=halogen atom



All the desired compound of formula (I) of the present invention can be synthesized according to the synthetic method (I) and, the compound of formula (I) wherein  $R^2$  is hydrogen can also be particularly prepared according to the synthetic method (II).

30 Hereinafter, the synthetic methods of the present invention will be more specifically explained.

#### Synthetic Method (I)

35 The compound of formula (I) according to the present invention can be synthesized by the following method.

When in the compound of formula (I)  $R^1$  represents alkyl, alkenyl, cycloalkyl, phenyl or benzyl and  $R^2$  represents hydrogen, or alkyl which is different from  $R^1$ , first, the intermediate compounds of formulae (4-1) and (4-2) are prepared according to the following reaction scheme (1).


40

45

50

55

## Reaction Scheme (1)



In the above reaction scheme,  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  are defined as previously described.

The precursor derivative of formula (1) which is used as the starting compound in the above reaction scheme (1) can be prepared from ammonium thiocyanate, benzoyl chloride and primary amine according to the method disclosed in O. S. Coll., Vol. 3, 734; and the precursor derivative of formula (2) which is also used as the starting material in the reaction scheme (1) can be prepared by reacting ethyl  $\beta$ -ketone ester either with sulfonyl chloride (see Synthesis, (1987), p188) or with chlorine.

The compound of formula (3-1) can be obtained by condensing the compound of formula (1) with the compound of formula (2) in the presence of a solvent. As the solvent for this purpose, ketones such as acetone, methylethyl ketone, etc., ethers such as tetrahydrofuran, diethylether, etc., halogenated hydrocarbons such as dichloromethane, 1,2-dichloroethane, chloroform, etc., alcohols such as methanol, ethanol, etc., and the like, can be preferably used. Among them, halogenated hydrocarbon solvents are most preferably used. The reaction can be carried out at the temperature in the range of 40°C to 120°C and most preferably at the temperature in the range of 70°C to 100°C.

The compound of formula (4-1) can be obtained by hydrolysis of the compound of formula (3-1) in the presence of a base. In this hydrolysis reaction, an inorganic base such as sodium hydroxide, potassium hydroxide, etc., is suitable as the base; and as the solvent a mixed solvent of ethers such as tetrahydrofuran, etc., and water or a mixed solvent of alcohols such as methanol, ethanol, etc., and water can be preferably used. This reaction can be practiced at the temperature in the range of 20°C to 120°C.


When  $\text{R}^1$  differs from  $\text{R}^2$ , the compound of formula (3-2) can be obtained by reacting the compound of formula (3-1) with various alkyl halides in the presence of a base. As the base which can be preferably used for this purpose, an organic base such as triethylamine, pyridine, etc., or an inorganic base such as sodium carbonate, sodium hydrogen carbonate, potassium carbonate, potassium hydride, sodium hydride, etc., can be mentioned, with sodium hydride being most effective. The solvent which can be used in this reaction includes halogenated hydrocarbons such as dichloromethane, 1,2-dichloroethane, etc., ethers such as tetrahydrofuran, diethylether, etc., amides such as dimethylformamide, dimethylacetamide, etc., and the like.

# EP 0 639 574 A1

The compound of formula (4-2) can be prepared by hydrolyzing the compound of formula (3-2) in the same manner as in the hydrolysis of the compound of formula (3-1).

When the compound of formula (I) wherein R<sup>1</sup> and R<sup>2</sup> represent the same alkyl group, or R<sup>1</sup> represents

5



10 and R<sup>2</sup> represents hydrogen is desired, the intermediate compounds of formulae (4-3) and (4-4) can be prepared according to the following reaction scheme (2).

15

20

25

30






35

40

45

50


55

Reaction Scheme (2)


The intermediate compound of formula (4-3) wherein  $\text{R}^1$  and  $\text{R}^2$  represent the same radical (R) can be prepared by preparing the compound of formula (3-1) having no substituent on the 2-amino group and reacting the compound of formula (3-1) thus prepared with 2 equivalent weights of the compound of formula RX wherein X represents halogen under the same condition as in the procedure for preparing the compound of formula (3-2) from the compound of formula (3-1) to prepare the compound of formula (3-3)

which is then hydrolyzed.


The intermediate compound of formula (4-4) wherein R' represents



can be prepared by reacting the compound of formula (3-1) having no substituent on the 2-amino group with an alkyl chloroformate




or an acyl halide



to obtain the compound of formula (3-4) which is then hydrolyzed.

Among the final desired compound of formula (I) according to the present invention, the compound of formula (7) can be prepared from the intermediate compound of formula (4) according to the following reaction scheme (3).

### Reaction Scheme (3)




The compound of formula (6) is prepared by stirring an aldehyde in ammonia water in which potassium cyanide and ammonium chloride are dissolved, according to Strecker synthesis (see O. S. Coll., Vol. 3, 84) which is one of the methods for synthesis of  $\alpha$ -aminonitriles.

The compound of formula (7) can be prepared by the following procedure. First, the compound of formula (4) is stirred together with thionyl chloride( $\text{SOCl}_2$ ) or phosphorus pentachloride( $\text{PCl}_5$ ) in a solvent, for example, halogenated alkyl hydrocarbons such as dichloromethane, chloroform, etc., aryl hydrocarbons such as benzene, toluene, xylene, etc., and the like, at the temperature in the range of  $0^\circ\text{C}$  to  $120^\circ\text{C}$  to obtain an acid halide of formula (5). The acid halide (5) thus obtained is then reacted with the compound of formula (6) in the presence of a base and a solvent to prepare the compound of formula (7). The base which can be suitably used in this reaction includes an organic base such as triethylamine, pyridine, etc., or an inorganic base such as sodium hydrogen carbonate, sodium carbonate, potassium carbonate, etc.; and as the solvent halogenated hydrocarbons such as dichloromethane, 1,2-dichloroethane, chloroform, carbon tetrachloride, etc., ethers such as tetrahydrofuran, diethylether, etc., ketones such as acetone, methylethyl ketone, etc., nitriles such as acetonitrile, isobutyronitrile, etc., and the like can be preferably used. This reaction can be carried out at the temperature in the range of  $0^\circ\text{C}$  to  $80^\circ\text{C}$ .

#### Synthetic Method (II)

The compound of formula (I) according to the present invention wherein  $\text{R}^1$  represents alkyl, alkenyl, cycloalkyl or benzyl and  $\text{R}^2$  represents hydrogen, i.e. the compound of formula (7-1), can be synthesized according to the following method.

#### Reaction Scheme (4)




Specifically, the compound of formula (7-1) can be obtained by reacting the compound of formula (8) with various primary amines of formula (9) in the presence of a base and a solvent. As the base suitable for this purpose, an organic base such as triethylamine, pyridine, etc., or an inorganic base such as sodium hydrogen carbonate, sodium carbonate, potassium carbonate, etc., can be used. In addition, when the primary amine which participates in the reaction is used in an excessive amount (2 to 3 times molar amount), this primary amine is used instead of the base. The most preferable base in this reaction is an organic base such as triethylamine. As the solvent halogenated hydrocarbons such as dichloromethane, 1,2-dichloroethane, chloroform, carbon tetrachloride, etc., ethers such as tetrahydrofuran, diethylether, etc., ketones such as acetone, methylethyl ketone, etc., amides such as dimethylformamide, dimethylacetamide, etc., and the like can be used, with ether solvents being most preferable.

In this reaction, the primary amine (9) is used, after any water is removed therefrom, in an amount of 2 to 3 moles with respect to one mole of the compound of formula (8). The reaction according to the reaction scheme (4) can be carried out at the temperature of  $20^\circ\text{C}$  to  $100^\circ\text{C}$ .

The derivative of formula (8) which is used as the starting compound in the above reaction can be synthesized by the method disclosed in European Patent No. 313,091, as depicted in the following reaction scheme (5).

Reaction Scheme (5)

5




Specifically, the compound of formula (8) is prepared by hydrolyzing the compound of formula (10), reacting the hydrolyzed product with thionyl chloride( $\text{SOCl}_2$ ) or phosphorus pentachloride( $\text{PCl}_5$ ) to obtain an acid halide of formula (11) which is then reacted with the  $\alpha$ -aminonitrile compound of formula (6).

30 The compound of formula (10) which is used as the starting compound for preparing the compound of formula (8) can be prepared by the method disclosed in J. Heterocyclic Chem., 22, 621 (1985).

The compound of formula (12) which is one of the final desired compound of the present invention can be prepared by the following reaction scheme (6).

35


Reaction Scheme (6)

50 Specifically, the compound of formula (12) can be prepared by stirring the compound of formula (7) in a base and a solvent while introducing hydrogen sulfide gas into the reaction solution. The suitable base in this reaction includes an organic base such as triethylamine, pyridine, etc., or an inorganic base such as sodium hydrogen carbonate, sodium carbonate, potassium carbonate, etc.; and as the solvent an organic base such as triethylamine, pyridine, etc., is used without any additional solvent or ethers such as tetrahydrofuran, diethylether, etc., halogenated hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, etc., esters such as ethylformate, ethyl acetate, etc., alcohols such as methanol, ethanol, etc., and the like can also be used. The reaction temperature can be  $20^\circ\text{C}$  to  $100^\circ\text{C}$ .

55

Typical compounds of formula (I) according to the present invention which can be prepared by the synthetic methods as described above are listed in the following Table 1.

Table 1









Comp. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
1	CH <sub>3</sub> -	H	CH <sub>3</sub> -		-CN
2	CH <sub>3</sub> -	CH <sub>3</sub> -	CH <sub>3</sub> -	"	-CN
3	CH <sub>3</sub> -	H	CH <sub>3</sub> -	"	$\begin{array}{c} \text{S} \\ \parallel \\ -\text{CNH}_2 \end{array}$
4	CH <sub>3</sub> -	H	CF <sub>3</sub> -	"	-CN
5	CH <sub>3</sub> CH <sub>2</sub> -	H	CH <sub>3</sub> -	"	-CN
6	CH <sub>3</sub> CH <sub>2</sub> -	H	CH <sub>3</sub> -	"	$\begin{array}{c} \text{S} \\ \parallel \\ -\text{CNH}_2 \end{array}$
7	CH <sub>3</sub> CH <sub>2</sub> -	H	CF <sub>3</sub> -	"	-CN
8	$\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{CH} - \\ \diagup \\ \text{CH}_3 \end{array}$	H	CH <sub>3</sub> -	"	-CN

Table 1 (continued)

5	Comp. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sub>4</sub>	R <sub>5</sub>
10	9	CH <sub>2</sub> =CH-CH <sub>2</sub> -	H	CH <sub>3</sub> -		-CN
15	10	CH <sub>2</sub> =CH-CH <sub>2</sub> -	H	CH <sub>3</sub> -	"	$\text{-}\overset{\text{S}}{\parallel}\text{CNH}_2$
20	11	CH <sub>2</sub> =CH-CH <sub>2</sub> -	H	CF <sub>3</sub> -	"	-CN
25	12		H	CH <sub>3</sub> -	"	"
30	13	"	H	CH <sub>3</sub> -		"
35	14	"	H	CH <sub>3</sub> -		$\text{-}\overset{\text{S}}{\parallel}\text{CNH}_2$
40	15	"	H	CF <sub>3</sub> -	"	-CN
45	16	"	H	CH <sub>3</sub> -		"
50	17	"	H	CH <sub>3</sub> -	-Ph	"

55

Table 1 (continued)













Comp. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
18	(CH <sub>3</sub> ) <sub>3</sub> C-	H	CH <sub>3</sub> -		-CN
19	PhCH <sub>2</sub> -	H	CH <sub>3</sub> -	"	"
20	"	H	CH <sub>3</sub> -		"
21	PhCH <sub>2</sub> -	H	CH <sub>3</sub> -	-Ph	"
22	Ph-	H	CH <sub>3</sub> -		"
23	"	H	CH <sub>3</sub> -		"
24	"	H	CH <sub>3</sub> -	-Ph	"
25	CH <sub>3</sub> OC(=O)-	H	CH <sub>3</sub> -		"
26	"	H	CH <sub>3</sub> -	"	

Table 1 (continued)

5	Comp. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
10	27	$\text{CH}_3\text{OC}(=\text{O})-$	H	$\text{CF}_3-$		-CN
15	28	"	H	$\text{CH}_3-$		"
20	29	"	H	$\text{CH}_3-$	-Ph	"
25	30	$\text{CH}_3\text{CH}_2\text{OC}(=\text{O})-$	H	$\text{CH}_3-$		"
30	31		H	$\text{CH}_3-$	-Ph	"
35	32	$(\text{CH}_3)_3\text{CC}(=\text{O})-$	H	$\text{CH}_3-$	"	"
40	33	$\text{PhC}(=\text{O})-$	H	$\text{CH}_3-$	"	"
45	34		H	$\text{CH}_3-$		"
50						

55

Table 1 (continued)
















Comp. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
35		H	CH <sub>3</sub> -	-Ph	-CN
36	CH <sub>3</sub> CH <sub>2</sub> -	H	CF <sub>3</sub> -		
37	CH <sub>3</sub> CH <sub>2</sub> -	H	CH <sub>3</sub> CH <sub>2</sub> -		-CN
38	CH <sub>3</sub> CH <sub>2</sub> -	H	CH <sub>3</sub> CH <sub>2</sub> -		
39	CH <sub>3</sub> -	H	CH <sub>3</sub> CH <sub>2</sub> -		-CN
40	CH <sub>3</sub> -	H	CH <sub>3</sub> CH <sub>2</sub> -		
41		H	CH <sub>3</sub> CH <sub>2</sub> -		-CN

Table 1 (continued)

Comp. No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
42		H	CH <sub>3</sub> CH <sub>2</sub> -		
43	CH <sub>2</sub> =CH-CH <sub>2</sub> -	H	CH <sub>3</sub> CH <sub>2</sub> -		-CN


The compound of formula (I) according to the present invention has a potent fungicidal activity against various phytopathogenic organisms, particularly against Oomycetes, and therefore, can be used as an agent for controlling plant diseases caused by such phytopathogenic organisms, for example, *Plasmopara viticola* in grapes, *Phytophthora infestans* in potato, *Phytophthora capsici* in red pepper, etc.

When the compound of formula (I) according to the present invention is used for controlling plant diseases in agricultural field, the compound of formula (I) is can be prepared in the form of a agricultural composition. Such agricultural composition contains at least one of the compound of formula (I) as an active ingredient, together with a conventional agriculturally acceptable carrier.

The present invention will be more specifically explained by the following preparations and examples. However, it should be understood that the present invention will not be limited to those examples in any manner.


#### PREPARATION 1

##### Synthesis of 2-methylamino-4-methyl-thiazole-5-carboxylic acid




3.6g of methyl thiourea and 7.9g of ethyl 2-chloroacetoacetate were added to 100ml of 1,2-dichloroethane and the mixture was stirred under refluxing for 10 hours. After removing the solvent under reduced pressure, 10% aqueous sodium hydroxide solution was added thereto to recrystallize the resulting product which was then filtered to obtain 7.8g (Yield 96%) of ethyl 2-methylamino-4-methyl-thiazole-5-carboxylate as a pale yellow solid. The obtained solid product was dissolved in 60ml of the mixed solvent of methanol-distilled water (v/v = 3/1) containing 1.8g of sodium hydroxide. The reaction mixture was stirred under refluxing for 8 hours, evaporated under reduced pressure to remove methanol and then adjusted to pH 2 to 3 by adding 10% aqueous hydrochloric acid solution to precipitate the white solid. The precipitated solid product was filtered, washed with water and diethylether and then dried to obtain 6.0g (Yield 90%) of the title compound as a white solid.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) : δ 12.4(1H, br), 3.0(3H, s), 2.40(3H, s)

PREPARATION 2Synthesis of 2-dimethylamino-4-methyl-thiazole-5-carboxylic acid


1.6g of ethyl 2-amino-4-methyl-thiazole-5-carboxylate and 0.68g of sodium hydride (purity 60%) were dissolved in 20ml of dry tetrahydrofuran and then 2.4ml of iodomethane was added thereto at 0°C. The reaction mixture was stirred for one hour at normal temperature and then extracted with water and ethylacetate. The extract was dried over anhydrous magnesium sulfate and evaporated to obtain 1.6g (Yield 89%) of ethyl 2-dimethylamino-4-methyl-thiazole-5-carboxylate as a white solid. The resulting product was added to 60ml of the mixed solvent of tetrahydrofuran-water (v/v = 2/1) containing 1.2g of sodium hydroxide. The reaction mixture was stirred for 16 hours at normal temperature and then treated according to the same procedure as PREPARATION 1 to obtain 1.3g (Yield 86%) of the title compound as a white solid.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) : δ 11.8(1H, br), 3.3(6H, s), 2.50(3H, s)

PREPARATION 3Synthesis of 2-methoxycarbonylamino-4-methyl-thiazole-5-carboxylic acid

2.0g of ethyl 2-amino-4-methyl-thiazole-5-carboxylate and 2.7ml of triethylamine were dissolved in 40ml of dichloromethane and then 1.1ml of methyl chloroformate added thereto at 0°C. The reaction mixture was stirred for 6 hours at normal temperature, evaporated to remove the solvent and then extracted with 10% aqueous sodium hydroxide solution and ethylacetate. The organic layer was separated, dried over anhydrous magnesium sulfate and then evaporated to obtain 2.1g (Yield 81%) of ethyl 2-methoxycarbonylamino-4-methyl-thiazole-5-carboxylate as a pale yellow solid. The resulting product was hydrolyzed according to the same procedure as PREPARATION 1 to obtain 1.4g (Yield 75%) of the title compound as a pale yellow solid.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) : δ 12.3(1H, s, br), 3.80(3H, s), 2.50(3H, s)

PREPARATION 4Synthesis of 2-bromo-4-trifluoromethyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide

15 1.2g of 2-bromo-4-trifluoromethyl-thiazole-5-carboxylic acid and 1.0g of phosphorus pentachloride were added to 10ml of dichloromethane and then stirred under refluxing for one hour. After removing the solvent and phosphorus oxychloride under reduced pressure, the residue was dissolved again in 10ml of dichloromethane and 0.84g of amino-thiophen-2-yl-acetonitrile hydrochloride and 1.3ml of triethylamine were added thereto at 0°C. The reaction mixture was stirred for 2 hours at normal temperature. Water was added

20 to the reaction mixture and the organic layer was separated. The separated organic layer was dried over anhydrous magnesium sulfate and then evaporated. The residue was then subjected to silica gel column chromatography to obtain 1.2g (Yield 70%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.42(1H, d), 7.35(1H, d), 7.06(1H, t), 6.40(1H, d, br), 6.25(1H, br)

25 PREPARATION 5Synthesis of 2-bromo-4-ethyl-thiazole-5-carboxylic acid (cyanothiophen-2-yl-methyl)-amide


40 According to the same procedure as PREPARATION 4 except that 1.5g of 2-bromo-4-ethyl-thiazole-5-carboxylic acid is used instead of 2-bromo-4-trifluoromethyl-thiazole-5-carboxylic acid, 1.68g (Yield 75%) of the title compound was obtained.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.41(1H, d), 7.34(1H, d), 7.05(1H, t), 6.40(2H, br)

45


50

55

EXAMPLE 1Synthesis of 2-methylamino-4-methyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (1)

1.8g of 2-methylamino-4-methyl-thiazole-5-carboxylic acid and 2.4g of phosphorus pentachloride were added to 20ml of dichloromethane and then stirred under refluxing for one hour. After removing the solvent and phosphorus oxychloride under reduced pressure, the residue was dissolved again in 20ml of dichloromethane and 2.0g of amino-thiophen-2-yl-acetonitrile hydrochloride and 4.8ml of triethylamine were added thereto 0°C. The reaction mixture was stirred for 2 hours at normal temperature and then the solvent was removed under reduced pressure. The residue was extracted with 10% aqueous sodium hydroxide solution and ethylacetate. The organic layer was separated, dried over anhydrous magnesium sulfate and then evaporated. The residue was then subjected to silica gel column chromatography and then the desired fraction was recrystallized from n-hexane and ethylacetate to obtain 1.05g (Yield 34%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.39(1H, d), 7.34(1H, d), 7.04(1H, t), 6.43(1H, d), 5.97(1H, d), 5.90(1H, s, br), 3.0-(3H, s), 2.55(3H, s)


EXAMPLE 2Synthesis of 2-dimethylamino-4-methyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (2)

1.2g of 2-dimethylamino-4-methyl-thiazole-5-carboxylic acid was converted into 2-dimethylamino-4-methyl-thiazole-5-carboxylic acid chloride using 1.3g of phosphorus pentachloride according to the same procedure as EXAMPLE 1. Then 1.1g of amino-thiophen-2-yl-acetonitrile hydrochloride and 2.5ml of triethylamine were added thereto and the reaction mixture was treated according to the same procedure as EXAMPLE 1 to obtain 0.65g (Yield 36%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.39(1H, d), 7.34(1H, d), 7.04(1H, t), 6.42(1H, d), 6.00(1H, d), 3.13(6H, s), 2.56-(3H, s)

**EXAMPLE 3**

Synthesis of 2-ethylamino-4-methyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (5)




1.2g of 2-ethylamino-4-methyl-thiazole-5-carboxylic acid was converted into 2-ethylamino-4-methyl-thiazole-5-carboxylic acid chloride using 1.3g of phosphorus pentachloride according to the same procedure as EXAMPLE 1. Then 1.1g of amino-thiophen-2-yl-acetonitrile hydrochloride and 2.5ml of triethylamine were added thereto and the reaction mixture was treated according to the same procedure as EXAMPLE 1 to obtain 0.57g (Yield 32%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.36(1H, d), 7.30(1H, d), 7.04(1H, t), 6.10(1H, d), 5.99(1H, s, br), 3.28(2H, q), 2.53(3H, s), 1.30(3H, t)

**EXAMPLE 4**

Synthesis of 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (12)




1.7g of 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid was converted into 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid chloride using 1.9g of phosphorus pentachloride according to the same procedure as EXAMPLE 1. Then 1.7g of amino-thiophen-2-yl-acetonitrile hydrochloride and 3.9ml of triethylamine were added thereto and the reaction mixture was treated according to the same procedure as EXAMPLE 1 to obtain 1.0g (Yield 39%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.40(1H, d), 7.34(1H, d), 7.05(1H, t), 6.60(1H, s), 6.45(1H, d), 6.00(1H, d), 2.59(1H, m), 2.54(3H, s), 0.80(4H, m)

**EXAMPLE 5**

Synthesis of 2-t-butylamino-4-methyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (18)




1.5g of 2-t-butylamino-4-methyl-thiazole-5-carboxylic acid was converted into 2-t-butylamino-4-methyl-thiazole-5-carboxylic acid chloride using 1.6g of phosphorus pentachloride according to the same procedure as EXAMPLE 1. Then 1.4g of amino-thiophen-2-yl-acetonitrile hydrochloride and 3.3ml of triethylamine were added thereto and the reaction mixture was treated according to the same procedure as EXAMPLE 1 to obtain 0.96g (Yield 41%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.38(1H, d), 7.34(1H, d), 7.04(1H, t), 6.44(1H, d), 6.00(1H, d), 5.56(1H, s, br), 2.52(3H, s), 1.42(9H, s)

**EXAMPLE 6**

Synthesis of 2-methoxycarbonylamino-4-methyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (25)



1.7g of 2-methoxycarbonylamino-4-methyl-thiazole-5-carboxylic acid was converted into 2-methoxycarbonylamino-4-methyl-thiazole-5-carboxylic acid chloride using 1.8g of phosphorus pentachloride according to the same procedure as EXAMPLE 1. Then 1.5g of amino-thiophen-2-yl-acetonitrile hydrochloride and 2.4ml of triethylamine were added thereto and the reaction mixture was treated according to the same procedure as EXAMPLE 1 to obtain 0.85g (Yield 32%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 10.5(1H, s), 7.41(1H, d), 7.35(1H, d), 7.06(1H, t), 6.41(1H, d), 6.18(1H, d), 3.89(3H, s), 2.66(3H, s)


The compound Nos. 8, 9, 19, 22 and 30 were synthesized according to the same procedure as EXAMPLES 1 to 6. <sup>1</sup>H-NMR data of the synthesized compounds are described in the following Table 2.

Table 2

Comp. No.	<sup>1</sup> H-NMR (solvent) : δ	
8	(CDCl <sub>3</sub> ):	7.38(1H, d), 7.30(1H, d), 7.00(1H, t), 6.44 (1H, d), 6.00(1H, d), 5.49(1H, s, br), 3.61 (1H, m), 2.52(3H, s), 1.27(6H, m)
9	(CDCl <sub>3</sub> ):	7.38(1H, d), 7.32(1H, d), 7.04(1H, t), 6.50 (1H, s, br), 6.42(1H, d), 6.01(1H, d), 5.85 (1H, m), 5.30(1H, d), 5.25(1H, m), 3.88(2H, m), 2.53(3H, s)
19	(CDCl <sub>3</sub> ):	7.35(7H, m), 7.04(1H, t), 6.42(1H, d), 6.25 (1H, s), 5.95(1H, d), 4.46(2H, s), 2.51(3H, s)
22	(CDCl <sub>3</sub> ):	8.40(1H, s), 7.35(6H, m), 7.20(1H, t), 7.00 (1H, s), 6.40(1H, d), 6.00(1H, d), 2.55(3H, s)
30	(CDCl <sub>3</sub> ):	10.70(1H, s, br), 7.37(1H, d), 7.30(1H, d), 7.05(1H, t), 6.39(1H, d), 6.20(1H, d), 4.30 (2H, q), 2.67(3H, s), 1.34(3H, t)

**EXAMPLE 7**

Synthesis of 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid (cyano-thiophen-3-yl-methyl)-amide (13)




1.4g of 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid was converted into 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid chloride using 1.6g of phosphorus pentachloride according to the same procedure as EXAMPLE 1. Then 1.4g of amino-thiophen-3-yl-acetonitrile hydrochloride and 3.2ml of triethylamine were added thereto and the reaction mixture was treated according to the same procedure as EXAMPLE 1 to obtain 0.73g (Yield 35%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.54(1H, s), 7.40(1H, m), 7.20(2H, m), 6.92(1H, s), 6.30(1H, d), 2.60(1H, m), 2.53-(3H, s), 0.80(4H, m)

**EXAMPLE 8**

Synthesis of 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid (cyano-furan-2-yl-methyl)-amide (16)



1.4g of 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid was converted into 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid chloride using 1.6g of phosphorus pentachloride according to the same

procedure as EXAMPLE 1. Then 1.2g of amino-furan-2-yl-acetonitrile hydrochloride and 3.2ml of triethylamine were added thereto and the reaction mixture was treated according to the same procedure as EXAMPLE 1 to obtain 0.83g (Yield 35%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.48(1H, d), 6.80(1H, s), 6.58(1H, d), 6.43(1H, t), 6.32(1H, d), 6.00(1H, d), 2.60-(1H, m), 2.55(3H, s), 0.85(4H, m)


The compound Nos. 20, 23, 28 and 34 were synthesized according to the same procedure as EXAMPLE 8. <sup>1</sup>H-NMR data of the synthesized compounds are described in the following Table 3.

Table 3

Comp. No.	<sup>1</sup> H-NMR (solvent) : δ	
20	(CDCl <sub>3</sub> ):	7.30(6H, m), 6.60(2H, m), 6.45(1H, s), 6.25 (1H, d), 5.95(1H, d), 4.45(2H, s), 2.50(3H, s)
23	(CDCl <sub>3</sub> ):	8.20(1H, s), 7.40(5H, m), 7.16(1H, m), 6.57 (1H, t), 6.42(1H, d), 6.30(1H, d), 6.00(1H, d), 2.57(3H, s)
28	(CDCl <sub>3</sub> ):	9.90(1H, s), 7.49(1H, s), 6.59(1H, d), 6.44 (1H, s), 6.27(1H, d), 6.15(1H, d), 3.89(3H, s), 2.64(3H, s)
34	(CDCl <sub>3</sub> ):	10.00(1H, s, br), 7.95(1H, d), 7.45(3H, m), 6.60(1H, s), 6.50(1H, s), 6.28(1H, d), 6.20 (1H, d), 2.65(3H, s)

#### EXAMPLE 9

Synthesis of 2-benzylamino-4-methyl-thiazole-5-carboxylic acid (cyano-phenyl-methyl)-amide (21)



1.2g of 2-benzylamino-4-methyl-thiazole-5-carboxylic acid was converted into 2-benzylamino-4-methyl-thiazole-5-carboxylic acid chloride using 1.1g of phosphorus pentachloride according to the same procedure as EXAMPLE 1. Then 0.89g of amino-phenylacetonitrile hydrochloride and 2.3ml of triethylamine were added thereto and the reaction mixture was treated according to the same procedure as EXAMPLE 1 to obtain 0.68g (Yield 39%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.40(10H, m), 6.25(1H, d), 6.15(1H, s, br), 5.80(1H, d), 4.45(2H, s), 2.52(3H, s)


The compound Nos. 17, 24, 29, 31, 32, 33 and 35 were synthesized according to the same procedure as EXAMPLE 9. <sup>1</sup>H-NMR data of the synthesized compounds are described in the following Table 4.

Table 4

Comp. No.	<sup>1</sup> H-NMR (solvent) : δ	
17	(CDCl <sub>3</sub> ):	7.46(5H, m), 7.28(1H, s), 6.29(1H, d), 5.95 (1H, d), 2.60(1H, m), 2.53(3H, s), 0.83(4H, m)
24	(CDCl <sub>3</sub> ):	8.0(1H, s, br), 7.30(9H, m), 7.15(1H, t), 6.25(1H, d), 5.85(1H, d), 2.55(3H, s)
29	(CDCl <sub>3</sub> ):	10.00(1H, s), 7.50(5H, m), 6.23(1H, d), 6.00(1H, d), 3.88(3H, s), 2.65(3H, s)
31	(CDCl <sub>3</sub> ):	9.40(1H, s), 7.50(5H, m), 6.20(1H, d), 6.00 (1H, d), 2.70(1H, m), 2.65(3H, s), 1.20(4H, m)
32	(CDCl <sub>3</sub> ):	8.90(1H, s), 7.50(5H, m), 6.20(1H, d), 6.00 (1H, d), 2.65(3H, s), 1.30(9H, s)
33	(CDCl <sub>3</sub> ):	7.90(2H, m), 7.65(9H, m), 6.20(1H, d), 6.15 (1H, d), 2.60(3H, s)
35	(CDCl <sub>3</sub> ):	10.1(1H, s, br), 7.90(1H, d), 7.50(7H, m), 6.25(1H, d), 6.15(1H, d), 2.60(3H, s)

**EXAMPLE 10**

Synthesis of 2-ethylamino-4-trifluoromethyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (7)




1.0g of 2-bromo-4-trifluoromethyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide, 0.36ml of triethylamine and 0.50ml of ethylamine were added to 10ml of tetrahydrofuran and the reaction mixture was stirred under refluxing for 4 hours. Water and ethylacetate were added thereto to separate the layers. The separated organic layer was dried over anhydrous sodium sulfate and evaporated. The residue was recrystallized from the mixed solvent of n-hexane and ethylacetate to obtain 0.67g (Yield 73%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.42(1H, d), 7.34(1H, d), 7.06(1H, t), 6.59(1H, s, br), 6.34(1H, d), 5.69(1H, s, br), 3.32(2H, q), 1.33(3H, t)


**EXAMPLE 11**

Synthesis of 2-ethylamino-4-ethyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (37)




1.0g of 2-bromo-4-ethyl-thiazole-5-carboxylic acid (cyanothiophen-2-yl-methyl)-amide, 0.40ml of triethylamine and 0.55ml of ethylamine were added to 10ml of tetrahydrofuran. The reaction mixture was stirred under refluxing for 4 hours and then treated according to the same procedure as EXAMPLE 10 to obtain 0.63g (Yield 70%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.39(1H, d), 7.33(1H, d), 7.05(1H, t), 5.90(1H, d), 5.50(1H, s, br), 3.28(2H, m), 2.92(2H, q), 1.28(6H, m)

**EXAMPLE 12****Synthesis of 2-methylamino-4-ethyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (39)**

1.0g of 2-bromo-4-ethyl-thiazole-5-carboxylic acid (cyanothiophen-2-yl-methyl)-amide, 0.40ml of triethylamine and 0.58ml of methylamine were added to 10ml of tetrahydrofuran. The reaction mixture was stirred under refluxing for 4 hours and then treated according to the same procedure as EXAMPLE 10 to obtain 0.77g (Yield 89%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.58(1H, d), 7.38(1H, d), 7.30(1H, d), 7.03(1H, t), 6.94(1H, s, br), 6.43(1H, d), 2.98(2H, q), 2.95(3H, s), 1.25(3H, t)

**EXAMPLE 13****Synthesis of 2-allylamino-4-ethyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (43)**

1.0g of 2-bromo-4-ethyl-thiazole-5-carboxylic acid (cyanothiophen-2-yl-methyl)-amide, 0.40ml of triethylamine and 0.66ml of allylamine were added to 10ml of tetrahydrofuran. The reaction mixture was stirred under refluxing for 12 hours and then treated according to the same procedure as EXAMPLE 10 to obtain 0.57g (Yield 61%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 7.36(1H, d), 7.30(1H, t), 6.95(1H, d), 6.50(1H, s, br), 6.43(1H, d), 5.86(1H, m), 5.29(1H, d), 5.25(1H, d), 3.90(2H, m), 2.99(2H, q), 1.25(3H, t)


The compound Nos. 11, 15 and 41 were synthesized according to the same procedure as EXAMPLES 10 to 13. <sup>1</sup>H-NMR data of the synthesized compounds are described in the following Table 5.

Table 5

Comp. No.	<sup>1</sup> H-NMR (solvent) : δ	
11	(CDCl <sub>3</sub> ):	7.41(1H, d), 7.33(1H, d), 7.05(1H, t), 6.60 (1H, s, br), 5.83(1H, m), 5.32(1H, d), 5.28 (1H, d), 3.90(2H, m)
15	(CDCl <sub>3</sub> ):	7.30(1H, d), 7.36(1H, d), 7.06(1H, t), 6.59 (1H, s, br), 6.38(1H, d), 6.28(1H, s), 2.64 (1H, m), 0.80(4H, m)
41	(CDCl <sub>3</sub> ):	7.75(1H, d), 7.36(1H, d), 7.31(1H, d), 7.09 (1H, s, br), 7.02(1H, t), 6.45(1H, d), 2.97 (2H, q), 2.56(1H, m), 1.24(3H, t), 0.72(4H, m)

## EXAMPLE 14

Synthesis of 2-ethylamino-4-methyl-thiazole-5-carboxylic acid (thiocarbamoyl-thiophen-2-yl-methyl)-amide (6)




0.85g of 2-ethylamino-4-methyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (5) was dissolved in 2ml of pyridine and the reaction solution was stirred for 6 hours at 60°C while introducing hydrogen sulfide gas. After removing pyridine, the residue was recrystallized from ethylacetate to obtain 0.86g (Yield 91%) of the title compound.

<sup>1</sup>H-NMR (Acetone-d<sub>6</sub>) : δ 9.25(2H, s, br), 7.56(1H, d), 7.35(1H, d), 7.22(1H, d), 7.14(1H, s), 6.95(1H, t), 6.26(1H, d), 3.33(2H, m), 2.49(3H, s), 1.23(3H, t)

## EXAMPLE 15

Synthesis of 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid (thiocarbamoyl-thiophen-2-yl-methyl)-amide (14)



0.72g of 2-cyclopropylamino-4-methyl-thiazole-5-carboxylic acid (cyano-thiophen-2-yl-methyl)-amide (12) was dissolved in 10ml of ethanol and then 0.63ml of triethylamine was added thereto. The reaction solution was stirred under refluxing for 8 hours while introducing hydrogen sulfide gas. After removing the solvent, water was added to the residue and then the solid product which is not dissolved in water was filtered. The filtered solid product was washed with diethylether and then dried to obtain 0.70g (Yield 88%) of the title compound.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 9.00(1H, s), 8.46(1H, s), 7.60(1H, d), 7.23(2H, m), 6.93(1H, t), 6.75(1H, s), 6.30(1H, d), 2.58(1H, m), 2.56(3H, s), 0.80(4H, m)

The compound Nos. 3, 10, 26, 36, 38, 40 and 42 were synthesized according to the same procedure as EXAMPLE 15. <sup>1</sup>H-NMR data of the synthesized compounds are described in the following Table 6.

Table 6

Comp. No.	<sup>1</sup> H-NMR (solvent) : δ
3	(Acetone-d <sub>6</sub> ): 9.20(2H, s, br), 7.55(1H, d), 7.34(1H, d), 7.33(1H, d), 7.22(1H, s, br), 6.94(1H, t), 6.25(1H, d), 2.94(3H, s), 2.49(3H, s)
10	(Acetone-d <sub>6</sub> ): 9.20(2H, s, br), 7.56(1H, d), 7.33(1H, d), 7.22(1H, d), 6.94(1H, t), 6.23(1H, d), 5.90(1H, m), 5.25(1H, d), 5.12(1H, d), 3.97 (2H, m), 2.49(3H, s)
26	(Acetone-d <sub>6</sub> ): 10.7(1H, s, br), 9.25(2H, s, br), 7.88 (1H, d), 7.37(1H, d), 7.26(1H, d), 6.96(1H, t), 6.27(1H, t), 3.82(3H, s), 2.60(3H, s)
36	(Acetone-d <sub>6</sub> ): 9.16(1H, s), 8.76(1H, s), 7.98(1H, d), 7.35(2H, m), 7.21(1H, m), 6.93(1H, s), 6.25 (1H, d), 3.26(2H, q), 1.26(3H, t)
38	(Acetone-d <sub>6</sub> ): 9.24(2H, d, br), 7.53(1H, d), 7.34(1H, d), 7.21(1H, d), 7.10(1H, s), 6.95(1H, t), 6.23(1H, d), 3.38(2H, m), 2.90(2H, q), 1.20 (6H, m)
40	(Acetone-d <sub>6</sub> ): 9.24(2H, d, br), 7.54(1H, d), 7.34(1H, d), 7.21(1H, d), 7.10(1H, s), 6.94(1H, t), 6.23(1H, d), 2.90(5H, m), 1.20(3H, t)
42	(Acetone-d <sub>6</sub> ): 9.24(2H, d, br), 7.65(2H, m), 7.36(2H, d), 7.25(2H, d), 6.97(1H, t), 6.28(1H, d), 2.94(2H, q), 2.62(1H, m), 1.22(3H, t), 0.74 (4H, m)

### Biological Test

#### Assay for determination of the fungicidal activity against phytopathogenic organisms


To identify the fungicidal effect of the compound of the present invention the activity against tomato late blight (*Phytophthora infestans*) and grape downy mildew (*Plasmopara viticola*) was determined.

#### Test 1

##### Preventive activity against tomato late blight

To determine the preventive activity the desired compound of the present invention was dissolved in 10% acetone solution and Tween-20 was added thereto in the concentration of 250ppm to prepare the test solution. The test solution was sprayed on tomato seedlings which were cultivated for 4 weeks in a greenhouse. Tomato seedlings on which the test solution was sprayed were allowed to stand for 24 hours at room temperature to evaporate the solvent and water. Then, the sporocyst suspension (10<sup>5</sup>/ml) of *Phytophthora infestans* which is the causative organism of tomato late blight was inoculated onto leaves of tomato. After 4 days, the state of tomato leaves was observed and compared with the state of the untreated group.

As the comparative drug dimethomorph having the following formula (VI) was used.



The state of tomato leaves was graded according to the following standard and described in the following Table 7.

EP 0 639 574 A1

Inhibition of development of plant disease	Grade
$\geq 90\%$	A
60-89%	B
$\leq 59\%$	C

5

10

15

20

25

30

35

40

45

50

55

Table 7

Concentration (ppm) Comp. No.	31	16	8	4
1	A	B	B	-
2	A	B	B	C
3	A	B	C	-
5	A	A	A	A
6	A	A	A	A
7	C	-	-	-
8	A	B	C	-
9	A	A	A	A
10	A	A	B	B
12	A	A	B	C
15	C	-	-	-
18	B	C	-	-
19	C	-	-	-
22	C	-	-	-
25	A	A	B	C
26	A	A	B	C
30	A	B	C	-
37	A	A	A	A
39	A	A	-	B
41	A	A	-	B
43	A	A	-	B
Dimethomorph	A	A	B	C

Test 2Systemic activity against tomato late blight

5 The test solution having a given concentration which was prepared according to the same method as in TEST 1 was injected in an amount of 5 ml into each pot (5cm X 5cm) containing tomato seedlings which were cultivated for approximately 4 weeks in a greenhouse. Then, the treated tomato seedlings were allowed to stand for 24 hours to evaporate the solvent. The sporocyst suspension ( $10^5$ /ml) of the causative organism of tomato late blight was inoculated on leaves of tomato. The inoculated tomato leaves were  
 10 maintained at 20°C and 100% relative humidity (RH) for 4 days to induce tomato late blight. Then, the development of tomato late blight was observed and compared with that in the untreated group to calculate the ratio of inhibiting the development of tomato late blight. The result is described in the following Table 8.

15 Table 8

20	Compound Concentration (ppm)	5	9	Dimethomorph
25	250	93	67	60
	63	58	0	25

30

TEST 3Curative activity against grape downy mildew

35 The curative activity of the compound of the present invention against grape downy mildew (*Plasmopara viticola*) was determined and then described in the following Table 9. The activity was graded A, B and C on the same standard as in TEST 1.


40 Table 9

45	Compound Concentration (ppm)	1	2	5	9	12	25
50	31	B	B	A	A	A	C

Although this invention has been described in its preferred form with a certain degree of particularity, it is appreciated by those skilled in the art that the present disclosure of the preferred form has been made  
 55 only by way of example and that numerous changes in the details of the construction, combination and arrangement of parts may be resorted to without departing from the spirit and scope of the invention.


## Claims

1. A novel 2-aminothiazolecarboxamide derivative represented by the following general formula (I) :



in which

15  $R^1$  and  $R^2$  independently of one another represent hydrogen,  $(C_1-C_5)$ alkyl,  $(C_1-C_5)$ haloalkyl,  $(C_3-C_6)$ alkenyl,  $(C_3-C_6)$ alkynyl,  $(C_3-C_6)$ cycloalkyl,



or phenyl or benzyl, each of which can be substituted with halogen,  $(C_1-C_3)$ alkyl or nitro;

25  $R^3$  represents  $(C_1-C_3)$ alkyl or  $(C_1-C_3)$ haloalkyl;

$R^4$  represents 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, phenyl, or phenyl or benzyl, each of which can be substituted with halogen,  $(C_1-C_6)$ alkyl or nitro;


$R^5$  represents cyano or thiocarbamoyl; and

$R^6$  represents  $(C_1-C_6)$ alkyl,  $(C_3-C_6)$ alkenyl,  $(C_3-C_6)$ cycloalkyl, or phenyl or benzyl, each of which can be substituted with halogen,  $(C_1-C_3)$ alkyl or nitro.

- 30 2. The compound of formula (I) according to claim 1 wherein the compound is in the form of an optically active isomer.

- 35 3. The compound of formula (I) according to claim 1 wherein

$R^1$  and  $R^2$  independently of one another represent hydrogen,  $(C_1-C_4)$ alkyl,  $(C_3-C_6)$ alkenyl,  $(C_3-C_6)$ alkynyl,  $(C_3-C_6)$ cycloalkyl or




45  $R^3$  represents methyl, ethyl or trifluoromethyl,

$R^4$  represents 2-thienyl or 3-thienyl,

$R^5$  represents cyano or thiocarbamoyl, and


$R^6$  represents  $(C_1-C_6)$ alkyl.

- 50 4. A process for preparing a compound having the following general formula (7) :



in which

$R^1$  and  $R^2$  independently of one another represent hydrogen,  $(C_1-C_5)$ alkyl,  $(C_1-C_5)$ haloalkyl,  $(C_3-C_6)$ alkenyl,  $(C_3-C_6)$ alkynyl,  $(C_3-C_6)$ cycloalkyl,




or phenyl or benzyl, each of which can be substituted with halogen,  $(C_1-C_3)$ alkyl or nitro;

$R^3$  represents  $(C_1-C_3)$ alkyl or  $(C_1-C_3)$ haloalkyl;


$R^4$  represents 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, phenyl, or phenyl or benzyl, each of which can be substituted with halogen,  $(C_1-C_6)$ alkyl or nitro; and

$R^6$  represents  $(C_1-C_6)$ alkyl,  $(C_3-C_6)$ alkenyl,  $(C_3-C_6)$ cycloalkyl, or phenyl or benzyl, each of which can be substituted with halogen,  $(C_1-C_3)$ alkyl or nitro,

characterized in that a compound having the following general formula (4) :



in which  $R^1$ ,  $R^2$  and  $R^3$  are defined as above, is reacted with thionyl chloride or phosphorus pentachloride to obtain an acid halide having the following general formula (5) :




in which  $R^1$ ,  $R^2$  and  $R^3$  are defined as above, and then the acid halide of formula (5) is reacted with a compound having the following general formula (6) :



in which  $R^4$  is defined as above.

5. A process for preparing a compound having the following general formula (7) :



in which


$R^1$  represents  $(C_1-C_5)$ alkyl,  $(C_1-C_5)$ haloalkyl,  $(C_3-C_5)$ alkenyl,  $(C_3-C_6)$ alkynyl,  $(C_3-C_6)$ cycloalkyl, or benzyl which can be substituted with halogen,  $(C_1-C_3)$ alkyl or nitro;

$R^2$  represents hydrogen;


$R^3$  represents  $(C_1-C_3)$ alkyl or  $(C_1-C_3)$ haloalkyl; and

$R^4$  represents 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, phenyl, or phenyl which can be substituted with halogen,  $(C_1-C_6)$ alkyl or nitro,

characterized in that a compound having the following general formula (8) :




in which  $R^3$  and  $R^4$  are defined as above and X represents halogen, is reacted with a primary amine having the following general formula (9) :




in which  $R^1$  is defined as above, in the presence of a base.

6. A process for preparing a compound having the following general formula (12) :



in which

$R^1$  and  $R^2$  independently of one another represent hydrogen,  $(C_1-C_5)$ alkyl,  $(C_1-C_5)$ haloalkyl,  $(C_3-C_6)$ alkenyl,  $(C_3-C_6)$ alkynyl,  $(C_3-C_6)$ cycloalkyl,




or phenyl or benzyl, each of which can be substituted with halogen,  $(C_1-C_3)$ alkyl or nitro;

$R^3$  represents  $(C_1-C_3)$ alkyl or  $(C_1-C_3)$ haloalkyl;

$R^4$  represents 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, phenyl, or phenyl or benzyl, each of which can be substituted with halogen,  $(C_1-C_6)$ alkyl or nitro; and

$R^6$  represents  $(C_1-C_6)$ alkyl,  $(C_3-C_6)$ alkenyl,  $(C_3-C_6)$ cycloalkyl, or phenyl or benzyl, each of which can be substituted with halogen,  $(C_1-C_3)$ alkyl or nitro,

characterized in that a compound having the following general formula (7) :



10 in which R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are defined as above, is reacted with hydrogen sulfide in the presence of a base and a solvent.

7. Use of the compound of formula (I) according to any one of claims 1 to 3 for controlling  
phytopathogenic organisms.

15 8. An agent for controlling phytopathogenic organisms which contain at least one of the compound of formula (I) as defined in claim 1.

20

25

30

35

40

45

50

55



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 94 11 2652

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	US-A-3 505 055 (VON SCHMELING ET AL.) 7 April 1970 * column 1; table VII * ---	1-8	C07D417/12 C07D277/56 A01N43/78
D,Y	EP-A-0 268 892 (NISSAN CHEMICAL INDUSTRIES LTD.) 1 June 1988 * claims 1,2,6-10 * ---	1-8	
D,Y	EP-A-0 292 937 (SUMITOMO CHEMICAL COMPANY LIMITED) 30 November 1988 * the whole document * ---	1-8	
Y	EP-A-0 296 673 (SHELL INTERNATIONALE RESEARCH MAATSCHAPPIJ B.V.) 28 December 1988 * claims 1-5,11-15 * ---	1-8	
D,Y	EP-A-0 313 091 (MITSUI TOATSU CHEMICALS INCORPORATED) 26 April 1989 * the whole document * ---	1-8	
D,Y	EP-A-0 434 620 (CIBA-GEIGY AG) 26 June 1991 * the whole document * -----	1-8	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 20 September 1994	Examiner Hartrampf, G
<b>CATEGORY OF CITED DOCUMENTS</b>			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	